

WRF input parameter updates based on recent and long-term satellite observations

Benas N., Chrysoulakis N., Christakis N., Kossioris G., Plexousakis M.

Recent satellite-derived products are analyzed and processed in order to update input parameters of the meso-scale model WRF (Weather Research and Forecasting), used for the study and prediction of extreme weather events in Greece, in the framework of the AKAIPRO¹ project. Updated input parameters include: the ASTER (Advanced Spaceborne Thermal Emission and Reflection Radiometer, onboard Terra satellite) derived GDEM (Global Digital Elevation Model), which substantially improves the original WRF elevation spatial resolution and surface emissivity and albedo derived from MODIS (Moderate Resolution Imaging Spectroradiometer, onboard Terra and Aqua satellites) multi-year observations. These emissivity and albedo timeseries, covering Europe, northern Africa and the Middle East, were used to update corresponding WRF input tables based on land cover classification after statistical processing. Updated input parameters also include land cover maps extracted from ESA's GlobCOVER product, based on MERIS (Medium Resolution Imaging Spectrometer, onboard Envisat platform) data. The processing and implementation procedures for updating the WRF input parameters are described in this study. WRF numerical simulations results and comparisons with data from meteorological stations are also presented, in order to assess the impact of the performed updates on the effectiveness of WRF in severe weather predictions.

Benas N.^{1*}, Chrysoulakis N.¹, Christakis N.^{2,3}, Kossioris G.³, Plexousakis M.³

1 Institute of Applied and Computational Mathematics, Foundation for Research and Technology Hellas.

2 Department of Physics, University of Crete, Greece.

3 Department of Mathematics, University of Crete, Greece.

*corresponding author e-mail: benas@physics.uoc.gr

¹ Study of extreme weather phenomena at a local scale and prediction of their impact on sectors of civil protection and the economy, Research Grant no. 09ΣΥΝ-31-1115, Program COOPERATION 2011, Ministry of Education and Religious Affairs, Greek Secretariat of Research Technology and EEYΔΕ-ETAK.

1 Introduction

Severe weather events are expected to change under changing climatic conditions, in terms of both their frequency and intensity. Among the most susceptible regions, the wider area of Greece is already witnessing such changes. The IPCC report on extreme events (IPCC 2012) concludes that there have been major increases in warm temperature extremes across the Mediterranean, while Trenberth (2011) found an increase in convective storm intensity over the same region. Extreme rainfall events are also increasing, according to e.g. Kioutsioukis et al. (2010).

Mesoscale numerical weather prediction models, such as the Weather Research and Forecasting (WRF) model, are essential for simulating and studying extreme events. In the AKAIPRO project, the WRF model is used for the study of extreme weather events in Greece during the last years, and their short-term prediction.

In the present study, recent satellite-derived products were analyzed and processed, in order to update important WRF input parameters, including topography, land cover, land surface emissivity (LSE) and land surface albedo (LSA). Topography, and especially its spatial resolution, is crucial for the accuracy of WRF simulations. Although increasing spatial resolution from 3km to 1km does not necessarily improve spatial verification statistics, higher resolution is needed to simulate correctly topography induced characteristics and structures (e.g. Mass et al. 2012). Land surface cover type variability, which determines the microclimate of a region but also affects local scale and mesoscale atmospheric circulation (e.g. Hartmann 1994), requires continuous updates. Updated LSE and LSA are needed for the computation of both shortwave and longwave radiation fluxes, which play a key role in surface-atmosphere interactions.

2 Data and Methodology

WRF represents topography using a Digital Elevation Model (DEM), derived from global observations of the Shuttle Radar Topography Mission (SRTM, Farr et al. 2007). The SRTM product is available at 900 m × 900 m spatial resolution, and was replaced in this study by the recently available Global DEM (GDEM), generated from data collected from the Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER), using photogrammetry and stereoscopy (Abrams et al. 2010). GDEM spatial resolution is 30 m × 30 m, having the potential to substantially improve the topography representation in the WRF.

The original WRF land cover data originate from the US Geological Survey (USGS) Global Land Cover Characteristics (GLCC) database, whereby land is divided in 24 categories, with a 900 m × 900 m spatial resolution. The data set was derived from AVHRR (Advanced Very High Resolution Radiometer) data, spanning from April 1992 through March 1993 (Loveland et al. 2000). These data were replaced by corresponding data from ESA's GlobCOVER product (2009), based on MERIS (Medium Resolution Imaging Spectrometer) data.

LSE and LSA in WRF are based on land cover types; two values of LSA (and similarly LSE) are assigned in each land cover category, for summer (16th April–15th October) and winter (16th October – 15th April). These data were derived from measurements by AVHRR (Csiszar and Gutman 1999), and were updated by data from MODIS, spanning the period 2000–2012.

2.1 Methodology for Topography, Land Cover, LSE and LSA Updates

For the GDEM incorporation in the WRF, a large number of GDEM tiles was initially retrieved, covering the wider Balkan and Central Mediterranean regions. These tiles were

used to create a mosaic image in GeoTIFF format, and in Geographic/WGS84 and EGM96 horizontal and vertical geodetic datums, respectively, as described in Chrysoulakis et al. (2011). The next step involved modification of this product into binary file format, compatible with the *geogrid* program of the WRF Preprocessing System (WPS), in order to be exploitable by the WRF, as well as changes in corresponding WPS input files. For this purpose, the methodology and software described in Beezley et al. (2011) were used. In the new topography, the GLCC land cover classification and topography interpolation method remained unchanged. Due to limitations in the number of topography input files used by the WRF, the spatial resolution of the GDEM data set was upscaled to $60 \text{ m} \times 60 \text{ m}$, using bilinear interpolation.

The number of land cover types differs between the USGS GLCC data set, used by the WRF, and GlobCOVER, which was used for updating these input data. Due to this difference, and in order to maintain the GLCC classification while updating its spatial distribution, land cover types of GlobCOVER were assigned to corresponding values of GLCC, based on their similarity.

In order to update the land cover types in the wider area of WRF simulations, 32 GLCC land cover tiles were converted from generic binary to GeoTIFF format and used for the creation of a mosaic image, which was then collocated with the corresponding GlobCOVER data. Furthermore, since the two data sets differ in their spatial resolution, ($900 \text{ m} \times 900 \text{ m}$ in GLCC and $300 \text{ m} \times 300 \text{ m}$ in GlobCOVER), necessitating a correspondence of 9 GlobCOVER pixels for each GLCC pixel, the updated value of each pixel was assigned based on the GlobCOVER land cover type which occupied the majority of these 9 pixels. Finally, the new data set was divided into 32 tiles, which replaced the initial GLCC WRF input.

WRF input LSE and LSA values were replaced by corresponding more recently available data from Terra MODIS Level 3, 8-day mean products, available at $1 \text{ km} \times 1 \text{ km}$ spatial resolution. For this purpose, 23 MODIS tiles, covering Europe, Northern Africa and the Middle East, were downloaded for each 8-day for the entire available time series (2000–2012), and processed accordingly (format conversion, data extraction and descaling).

LSE was initially calculated on a pixel basis, as the average of LSE in MODIS bands 31 and 32. Temporally averaged LSE values were then computed for the entire area, corresponding to summer and winter means. For the assignment of summer/winter value in each WRF land cover type, these seasonal mean LSE data were collocated with the corresponding land cover spatial distribution data, and the average LSE value for each land cover type was calculated.

LSA values were computed as a linear combination of black and white sky albedo values (BSA and WSA, respectively), as described in Schaaf et al. (2002). Specifically, 8-day mean BSA, which is a function of solar zenith angle (SZA), was first calculated on a pixel basis by averaging corresponding BSA values calculated at 9:00, 12:00, 15:00 and 18:00 local time SZAs, according to equation (2) in Schaaf et al. (2002). LSA was then calculated based on equation (3) in Schaaf et al. (2002), whereby it is a function of the fraction of diffuse skylight, which, in turn, is a function of aerosol optical thickness (AOT). For this purpose, AOT data from the corresponding MODIS Level 3 8-day product were used, covering the entire region, and LSA was computed by averaging results acquired using different SZAs, as previously described. The assignment of LSA values in each land cover type was performed with the same procedure described for LSE.

3 Case Study

The case study focused on a heat wave event (Metaxas and Kallos 1980) in Athens, in July 15th, 2012. Two WRF runs were performed, both with the new GDEM, and with the old and new land cover data, respectively. To represent the thermal and dynamic effects of urban surfaces, a single-layer Urban Canopy Model (UCM, Kusaka et al. 2001) was coupled in Noah Land Surface Model (LSM, Chen and Dudhia 2001).

Figure 1 shows the differences in the obtained WRF results for near surface air temperature, evaluated with the two land covers, at 2012-07-15 and 23:00 local time. The simulation based on the new land cover gave higher temperatures (by 0-5°C) in the northern and southern peri-urban areas of Athens, as well as in the eastern Attica (Mesogeia area), which was affected by substantial urban sprawl during the last decades (Chrysoulakis et al. 2013).

Two indicative comparative time series plots for selected stations are also presented. It should be noted, however, that comparisons with surface point measurements at specific locations can be misleading, given the specific local environment characteristics. In Fig. 2 (left, Gkazi station of the National Observatory of Athens), the new land cover slightly improves the simulated temperature. At Ano Liosia station, operated by the Hydrological Observatory of Athens (Fig. 2 right), the WRF simulation with the new land cover overestimates air temperature by 8.5% during the night of 15-07-2012. It follows, however, the exact temperature pattern during the night of 16-07-2012. On the other hand, the simulation with the old land cover underestimates by the same order of magnitude the temperatures during the nights of 15 and 16 -07-2012. Similar comparisons with other stations showed minor differences when the land cover type remained unchanged between the two data sets (relative error of the order of 5% or less). Moreover, daytime temperatures for all stations were underpredicted by the numerical model (for both land cover data sets) by 2-3°C on average, a fact which is in agreement with previous studies (Roux et al. 2008).




Fig. 1. Spatial distribution of the difference in near surface temperature (2m a.g.l.), between the two WRF runs, with the new and the old land cover, in July 15th, 2012, 23:00 local time.




Fig. 2. Intercomparison of the near surface temperature between the two WRF runs and Gkazi (left) and Ano Liosia (right) stations.

4 Conclusions

Important input parameters of the meso-scale WRF model were updated based on recently derived satellite products, in the framework of the AKAI PRO project, aiming to study

extreme weather events in Greece. These parameters included topography, land cover types distribution, LSE and LSA.

ASTER GDEM data were used for the topography update, improving its spatial resolution by two orders of magnitude. Recent land cover data from ESA's GlobCOVER product were also used to update the original, outdated WRF data set. For the LSE and LSA, MODIS data were used, covering a wide area (including Europe, Northern Africa and the Middle East), and spanning the period 2000-2012.

WRF simulations with the new land cover capture correctly the tendency for a stronger Urban Heat Island (UHI) effect in the Athens Greater Area, where extensive urbanization took place in the last decade (e.g. Giannaros et al. 2013). However, comparison of the simulated surface temperature with surface measurements at specific locations can be misleading given the specific local environment the measuring stations operate in. Further investigations are already under way, in order to assess these issues.

These preliminary results highlight the importance of up to date input parameters in WRF simulations, concerning regional climatic and environmental changes, which can lead to significant ecological and social consequences.

References

- Abrams M, Bailey B, Tsu H, Hato M (2010) The ASTER Global DEM. *Photogramm Eng Remote Sens* 76:344–348.
- Beezley JD, Kochanski AK, Mandel J (2011) Integrating high-resolution static data into WRF for real fire simulations. In: Proc. of the 9th Symposium on Fire and Forest Meteorology, Palm Springs, American Meteorological Society, Paper 6.3.
- Chen F, Dudhia J (2001) Coupling an Advanced Land Surface–Hydrology Model with the Penn State–NCAR MM5 Modeling System. Part I: Model Implementation and Sensitivity. *Mon Wea Rev*, 129:569–585.
- Chrysoulakis N, Mitraka Z, Stathopoulou M, Cartalis C (2013) A comparative analysis of the urban web of the greater Athens agglomeration for the last 20 years period on the basis of Landsat imagery. *Fresen Environ Bull* 22:2139–2144.
- Chrysoulakis N, Abrams M, Kamarianakis Y, Stanisławski M (2011) Validation of the ASTER GDEM for the area of Greece. *Photogramm Eng Remote Sens* 77:157–165.
- Csiszar I, Gutman G (1999) Mapping global land surface albedo from NOAA AVHRR. *J Geophys Res* 104 (D6):6215–6228.
- Farr TG, Rosen PA, Caro E, Crippen R, Duren R, Hensley S, Kobrick M, Paller M, et al. (2007) The Shuttle Radar Topography Mission. *Rev Geophys* 45(2). doi: 10.1029/2005RG000183
- IPCC (2012) *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptations*. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- Kioutsoukis I, Melas D, Zerefos C (2010) Statistical assessment of changes in climate extremes over Greece (1955–2002). *Int J Climatol* 30:1723–1737.
- Kusaka H, Kondo H, Kikegawa Y, Kimura F (2001) A simple single-layer urban canopy model for atmospheric models: Comparison with multi-layer and slab models. *Bound Layer Meteor*, 101:329–358.
- Loveland TR, Reed BC, Brown JF, Ohlen DO, Zhu J, Yang L, Merchant JW (2000) Development of a global land cover characteristics database and IGBP DISCover from 1-km AVHRR data. *Int J Remote Sens* 21:1303–1330.
- Mass CF, Ovens D, Westrick K, Colle BA (2002) Does increasing horizontal resolution produce more skillful forecasts? *Bull Amer Meteor Soc*, 83:407–430.
- Metaxas DA, Kallos G (1980) Heat waves from a synoptic point of view. *Riv Meteor Aeronaut* 2–3:107–119.
- Roux G, Liu Y, Monache LD, Shey RS, Warner TT (2008) Verification of High Resolution WRF-STFDDA Surface Forecasts over mountains and plains. 9th WRF workshop, Boulder, Colorado, 23-27 June 2008.
- Schaaf C, Gao F, Strahler A, Lucht W, Li X, Tsung T, Strugnell N, Zhang X, Jin Y, Muller JP, Lewis P, Barnsley M, Hobson P, Disney M, Roberts G, Dunderdale M, Doll C, d'Entremont R, Hu B, Liang S, Previtte J, Roy D (2002) First operational BRDF, albedo and nadir reflectance products from MODIS. *Remote Sens Environ* 83:135–148.
- Trenberth KE (2011) Changes in precipitation with climate change. *Clim Res* 47:123–138.

